

THE DECREASE IN NUMBER OF BLOOD POLYMORPHONUCLEAR (PMN) TO PERIAPICAL RADIOGRAPHS DOSE OF RADIATION EXPOSURE

PENURUNAN JUMLAH POLYMORPHONUCLEAR (PMN) DARAH TEPI PADA PAPARAN RADIASI SINAR X DOSIS RADIOGRAFI PERIAPIKAL

Amni Adlina¹, Wasilah²

¹ Fakultas Kedokteran Gigi Universitas Jember, Jember, Indonesia

² Staf Pengajar Kedokteran Gigi Institut Ilmu Kesehatan, Kediri, Indonesia

Korespondensi: Wasilah, email: wasilahyahya@gmail.com, Kedokteran Gigi Institut Ilmu Kesehatan, Jl. K.H. Wahid Hasyim 65, Kediri 64114, Jawa Timur. Telp. (0354) 773299, Faks (0354) 771539

Abstract

Background: Periapical radiograph is a type of radiographic examinations that is widely used in the field of dentistry. Radiographic X-ray is an ionizing radiation that can cause damage and death to cells, tissues or organs, including the polymorphonuclear (PMN). PMN is cell that serves as the first line of defense against the invasion of organisms. **Objective:** The purpose of this study is 1) to determine whether there is a decrease in the number of peripheral blood PMN after being given periapical radiograph dose of X-ray radiation exposure, and 2) to determine whether there is any difference in the declining number of peripheral blood PMN after being given a single dose and some different total dose of periapical radiograph of X-ray radiation exposure. **Method:** The design of this study is an experimental laboratory research with total sample of 24 strains of *Balb c* male mice. The sample is divided into four groups, each consisting 6 mice. Group 1 serves as the control group; Group 2 is given a treatment with a single-dose exposure of periapical radiographs of X-ray radiation; Group 3 is given a treatment of 6 times total doses exposure of periapical radiograph of X-ray radiation; and Group 4 is given a treatment of 14 times total doses exposure of periapical radiograph of X-ray radiation. After 24 hours of radiation exposure, the blood was drawn. **Conclusion :** The calculation is done by multiplying the number of PMN percentage from leukocyte counts with a total leukocyte. The data obtained are statistically tested using One Way Anova and LSD with a significance level of 95%. **Result :** The result of this study indicates that there is a difference in the peripheral blood PMN counts in male mice after being exposed to radiation doses of periapical radiographs in all groups ($P < 0.05$). There is an increasing number of peripheral blood PMN in male mice after being treated with X-ray radiation at a single dose of periapical radiographic exposure, whereas after being exposed with radiation at the dose of 6 and 14, the number of PMN is declining. The highest decreasing number of PMN is found in the group with exposure of radiation at 14 dose.

Keywords: Polymorphonuclear (PMN), X-ray Radiation, Periapical Radiograph

Introduction

Radiographic examination is a routine examination in the field of dentistry, thus radiography is a necessity that cannot be

avoided, and it can be said to be a very important thing for us ¹. Periapical radiograph is a type of radiographic examinations that is widely used in the field of dentistry ².

Periapical radiograph using X-ray, which is one of ionizing radiations³.

Lab examination found that few days of 25 rem (0,25 ceivert) X-ray radiation dosage influences blood condition i.e. anemia, leukopenia, thrombocytopenia, and leukemia. 0,5 gray (50 rad) dosage of ionizing radiation decreases neutrophil count in blood cells⁴. The primary function of neutrophil Polymorfonuklear (PMN) is to give imune response by performing phagocytosis and also killing or eradicating incoming microorganism⁵. In low dose radiation, few hours after an exposure the neutrophil count increases. However, after 24 hours and more of exposure neutrophil and leukocyte count is decreasing⁶.

There are some literatures that support the reduction in number of blood cells. Cole said that ionizing radiation can cause leukopenia. While Miller and Weller state that the ionizing radiation can lead to a reduction of all types of leukocytes. WHO also stated that ionizing radiation can cause an early haematological changes, but until now the decreased polymorphonuclear (PMN) due to periapical radiographs dose of X-ray exposure has not been studied⁷.

The purpose of this study is to determine whether there is a decrease in the number of peripheral blood PMN after being given periapical radiograph dose of X-ray radiation exposure, and to determine whether there is any difference in the declining number of peripheral blood PMN after being given a single dose and some different total dose of periapical radiograph of X-ray radiation exposure.

Materials and Methods

This study is a kind of experimental laboratory research. The study population is a strain of mice *Balb c.* male mice.

Grouping of samples stage

The control group (group 1), consisted of 6 healthy mice and not treated. Treatment group (group 2), consisted of 6 healthy mice into treatment group given a one-time exposure of dental radiodiagnostic. Treatment group

(group 3), consisted of 6 healthy mice into treatment groups that provided in 6 times dental radiodiagnostic exposure with an interval of time between them is 1 minute. Treatment group (group 4), consisted of 6 healthy mice into treatment groups that provided in 14 times dental radiodiagnostic exposure with an interval of time between them is 1 minute.

Mice fixation stage

The used fixation device is made of white plastic paralon. Bottom of the tube was perforated for the head of mice and six holes on the other edge was made by a stick. The mouse's head inserted through the hole on the bottom tube and the back is fixed with three crossing sticks between the two holes that made before.

Radiation exposure stages

Animal that has been fixed in white plastic paralon on a table in supine position (the abdomen is facing up). Cone of X-ray unit is directed right to the heart with the tip of the cone touches the surface of the plastic paralon. The animal is ready for the radiation exposure.

Collection and countin blood sample stage

Animals were sacrificed after 24 hours of exposure to radiation and the blood was taken from the heart using a 1 ml insulin syringe. The number of PMN was counted by multiplying the percentage of PMN in the leukocyte counts in blood smears by the number of leukocytes per mm³ in the counting room *Improver Neubeur*.

Statistical analysis

The obtained data were analyzed using the *Kolmogrof-Smirnof* normality test and the homogeneity test of *Levene*, followed by One Way ANOVA parametric test to determine whether there are groups of data that have significant mean differences, with 95% degree of significance ($p < 0.05$). Followed by LSD test (Least Significant Difference Test) to determine which groups were significantly different.

Result

The result of the study shows that the group with single dental radiodiagnostic exposure the absolute neutrophil count increases as much as 291.816,67. Meanwhile, the group which is given 6 times of dental radiodiagnostic exposure treatment shows that there is a decrease of absolute neutrophil count as much as 260.733,33. The number of absolute neutrophil count in the group that has received a treatment of 14 times exposure is the least one which is 236.933,33 and 260.733,33 (Table 1).

Table 1. Mean and standart deviation of the number of PMN in 24 hours observations post exposure of periapical radiographs dose of X-ray radiation

Treated and Control	Mean
Control group	280,333.33
P1	291,816.67
P2	260,733.33
P3	236,933.33

(cell/mm³)

P1 = treated group with single dose

P2 = treated group with a 6 times of exposure in total dose

P3 = treated group with a 14 times of exposure in total dose

Graph of the average number of PMN is shown in figure 1.




Fig.1. Histogram of the average number of PMN in post 24 hours observation

Before the data was analyzed, it was tested for normality and homogeneity first. The normality test used *Kolmogorof-Smirnof*, whereas the homogeneity test used *Levene test*. Both use the error rate (α) 0.05. In the used homogeneity test, $P = 0.974$ values obtained and the normality test, P value obtained for the four groups of data were > 0.05 .

Table 2. One Way Anova test results on the

	Sum of Square	df	Mean Square	F	Sig.
Between groups	10416571250.0	3	3472190417	3.1	.049
Within groups	22309288333.3	20	1115464417	13	
Total	32725859583.3	23			

control group and treatment

Anova test showed there were some differences in number of Neutrophyle. To find out which groups that significantly different, the data were analyzed using the LSD test which is shown in Table 3.

Table 3. LSD test result on four group

Multiple Comparisons

Dependent Variable: The number of neutrophil LSD

(I)	Treated group	Mean difference	Std. Error	Sig.
Control	P1	-	19282.67	.55
	P2	11483.333	3	8
	P3	19600.000	3	.32
		43400.000*	19282.67	2
P1	P2	31083.333	3	.03
		54883.333*	19282.67	.32
	P3		3	.01
P2	P3	23800.000	3	.06
			3	.23

*.The mean difference is significant at the ,05 level.

From the data analysis of LSD test to each group – control and experimental, shown on

Table 4.5 it can be seen that there is a significant difference between the control group and P14 group, between P1 group and P14 group. However, from the same data set, there is no significant difference between control group and P1 group; control group and P6 group; and P1 and P6 group.

Discussion

X-ray radiation not only providing benefits for all of us but also has a negative effect. Ionizing radiation is one of powerful agent in causing damage and even death of cells, tissues or organs, while the present technology cannot fully protect the body yet from the side effects of ionizing radiation⁴.

X-ray radiation from dental radiography units that used in this research is direct through a cone which has 100 cm of surface area. Cone is directed right to the heart of mice as the focus of the organ where blood will be taken, then they fixed so that cannot move and the radiation can certainly lead to cardiac of mice. Blood sampling performed on the heart with blood volume needed considerations for research and focus of the radiation had to be directed to specific organs, so that the heart was chosen as a spot of blood sampling. In this study used mice aged 3 to 4 months, because according to Schalm, mice in that age is not easy to die when irradiated below the lethal dose. Mice lethal dose ranged from 550-640 rad⁸.

The result of the research and the statistical test shows that there is a decrease in the number of peripheral blood count after given periapical radiograph of X-ray radiation exposure. This is also in line with Underwood's statement saying that radiation causes neutrophil reduction in blood circulation (opposite circulating blood?) alongside with other blood cells (pancytopenia)⁹. Edward also states that radiation has a negative effect to blood cells as it cuts down the number of cells in peripheral circulation^[4].

The decrease in the number of PMN neutrophil in this research is caused by a condition in which the effect of radiation exposure can be accumulated. This accords

Miles' statement saying that tissues have an ability to repair its damages after being exposed to radiation; however, unreparable damages can be accumulated¹⁰.

Cells damages can be caused both by direct and indirect effect of ionizing radiation. Damages due to direct effect happens when ionizing particles interact (energy transfer to) with biology macromolecule such as DNA, RNA, protein, or enzyme. Damages due to indirect effect happens when the damage is caused by oxidant agent resulted from ionizing process. 75% of cells is water and this water molecule is the one that is most ionized by X-rays¹⁰.

Result shows that there is an increase in PMN neutrophil count to the group with single exposure of dental radiodiagnostic of periapical radiation dose; however this increase is not so different with that of control group. This can be caused by the fact that oxidant of OH¹, H¹ and HOO¹ in blood circulation resulted from ionizing process is one of the inflammatory mediator that is transported to bone marrow capillaries to move the stored neutrophil into circulation blood⁴.

This research was conducted on male mice. Whether the results of this study can be equated or generalized to humans, of course, this requires a separate research and discussion, but please note that the lethal dosage (LD) 50/30 of mice is 550-640 rad¹¹. While the LD 50/30 of adult human is 450 rem (~450 rad). LD 50/30 is a whole body radiation dose that is lethal in 50% of the population within 30 days⁴.

Conclusions

1. There is a difference of peripheral neutrophil count in male mice *Balb c* after given an exposure of dental radiodiagnostic of periapical radiography between control group, single exposure group, 6x exposure group, and 14x exposure group.
2. There is an increase in peripheral neutrophil count in male mice *Balb c* after given a single exposure of X-ray radiation of periapical radiography dose; meanwhile after given 6x

and 14x of exposure, the number of PMN count is decreasing. The highest number of decrease in peripheral neutrophil count happens to the group with 14x of exposure.

References

1. Yunus, B. 2005: 'Radiografi Dental sebagai Diagnosa Awal untuk Mencegah Keparahan suatu Penyakit Gigi dan Mulut', *Majalah Kedokteran Gigi Edisi Khusus Temu Ilmiah Nasional*, 4(11) pp: 396-400
2. Suharjo And Endang. 1994: 'Peranan Teknik dan Interpretasi Radiografi Intraoral Periapikal dalam Perawatan Endodontic', *Jurnal Kedokteran Gigi PDGI*, 2(43) pp: 40-43
3. Cotran, R., Robbins, S., Kumar, Abbas and Nelson. 1999: 'Pathologic Basis of Disease', (Ed): 7, pp: 67-69
4. Edwards, Cris, Statkiewicz, M.A. and Russell, E. 1990. 'Radiation Protection for Dental Radiographers', pp: 43-48
5. Kresno, S.B. 2001: 'Imunologi: *Diagnosis dan Prosedur Laboratorium*', (Ed): 7, pp: 30
6. Rosyid, A. 2006. '*Temuan-Temuan pada Hasil Pemeriksaan Laboratorium akibat Radiasi*', pp: 24
7. Astuti, E.R. 1995: 'Pengaruh Radiasi Pengion dan Banyaknya Ulangan terhadap Total Leukosit, Hitung Jenis Leukosit, Hb, serta Total Eritrosit Mencit Jantan ', pp: 52-60
8. Goaz, P.W. and White, S.C. 1987: 'Oral Radiology', (Ed): 2, pp: 104-107
9. Underwood, J. C. E. 2000. *General and Systematic Pathology 3rd ed.* Edinburgh: Churchill Livingstone
10. Miles, Van Dis, Jensen, and Ferretti. 1993. *Radiographic Imaging for Dental Auxillaries*, (Ed): 2, pp: 34
11. Goaz, P.W. and White, S.C. 1987: 'Oral Radiology', (Ed): 2, pp: 104-107.